



# **Exact Draws from the Stationary Distribution of Entry-Exit Models**

Takashi Kamihigashi  
Research Institute for Economics and Business  
Administration, Kobe University

John Stachurski  
Research School of Economics, Australian National  
University

ANU Working Papers in Economics and Econometrics  
# 588

October 2012

JEL: C61, C63

ISBN: 0 86831 588 5

# Exact Draws from the Stationary Distribution of Entry-Exit Models\*

Takashi Kamihigashi<sup>a</sup> and John Stachurski<sup>b</sup>

<sup>a</sup>*Research Institute for Economics and Business Administration, Kobe University*

<sup>b</sup>*Research School of Economics, Australian National University*

October 17, 2012

## Abstract

---

In equilibrium models of firm dynamics, the stationary equilibrium distribution of firms summarizes the predictions of the model for a given set of primitives. Focusing on Hopenhayn's seminal model of firm dynamics with entry and exit (*Econometrica*, 60:5, 1992, p. 1127–1150), we provide an algorithm that generates *exact* draws from the stationary distribution in finite time for any specified exit threshold. The technique is able to rapidly generate large numbers of exact and independent draws.

*Keywords:* Simulation, stationary equilibrium, firm dynamics

*Journal of Economic Literature Classifications:* C61, C63

---

## 1 Introduction

In this paper we provide a method for generating exact, IID draws from the stationary distribution of Hopenhayn's (1992) model of firm dynamics with entry and exit. Hopenhayn's work is a cornerstone of the modern theory of firm dynamics, and forms the foundations of a large and growing literature (see, e.g., Hopenhayn and Rogerson, 1993; Cooley and Quadrini, 2001; Melitz, 2003 or Clementi and Palazzo, 2010). The stationary distribution of the model represents a cross-sectional distribution of firms that is both consistent with the definition of equilibrium at any point in time and also invariant over

---

\*We acknowledge financial support from ARC Discovery Outstanding Researcher Award DP120100321.

time. For typical parameter values the stationary distribution is uniquely defined but has no analytical solution. To compare the model with data, it is necessary to either compute or simulate from the distribution.

Direct computation of the distribution involves discretization of some form. The size of the resulting errors are typically unknown, reducing confidence in quantitative results. Regarding simulation, for a given exit policy it is straightforward to set up an ergodic Markov process such that the stationary distribution of the Markov process coincides with the cross-sectional stationary distribution of Hopenhayn's model, and this permits approximate simulation from the stationary distribution simply by running the process from an arbitrary initial value until it is judged to have "nearly" converged. Simulating until the distribution of the state is approximately stationary is referred to as "burn-in." Unfortunately, the length of burn-in required is the subject of guesswork and heuristics, which is clearly a poor foundation for analysis. Moreover, regardless of how much burn-in is performed, the resulting sample is never exactly stationary, and the size of the error is once again unknown. Finally, for a given method, the size of the error is likely to vary with the parameters, since the parameters change the structure of the problem. If the burn-in is not varied accordingly, this is likely to cause bias.

In this paper we show that these problems can in fact be overcome. By using a variation of Propp and Wilson's (1996) coupling from the past (CFTP) technique, we show that it is possible to generate *exact* independent draws from the stationary distribution of this class of models, for any specified exit threshold. An additional and related benefit of the algorithm is that it terminates as soon as an exact draw has been generated. Hence there is no need for the heuristics used to judge burn-in time. Moreover, by repeating the algorithm with independent seeds it becomes possible to generate multiple independent draws from the stationary distribution. These processes can be run sequentially or in parallel. Even with sequential generation, the algorithm permits rapid generation of exact IID draws from the stationary distribution. Because the resulting sample is IID, the standard central limit theorem for independent observations can be used to give confidence intervals for moments or empirical distributions.

The CFTP method has been used successfully for models with large but discrete state spaces, where standard methods for computing stationary distributions are infeasible. Applications range from statistical mechanics to page ranking and the design of peer-to-peer file sharing systems (see chapter 2 of Levin *et al.*, 2008 for recent survey). On the other hand, in the case of *continuous* state spaces, applicability of CFTP is not automatic. It depends on the particular structure of the model in question, and the ingenuity required to exploit it. Murdoch and Green (1998) showed that CFTP can in principle be used in continuous state settings when the underlying process satisfies Doeblin's condition. However, their method cannot necessarily be implemented in practice without approximating the model, thereby negating the exactness of the draws. Murdoch and

Green’s work was extended by Foss and Tweedie (1998), Athreya and Stenflo (2003) and many others. In particular, it is now well-known that CFTP can be highly efficient for monotone processes. Corcoran and Tweedie (2001) provide a general theory. Nishimura and Stachurski (2010) applied their ideas to an incomplete markets model.

The technique presented in this paper is entirely new. The state space is continuous, and, while the Markov process for productivity of incumbents is monotone increasing, the overall process that combines dynamics of incumbents and new entrants is not. Despite this, we develop a CFTP method by exploiting both the monotonicity of productivity for incumbents and the renewal property introduced by new entrants. It requires no approximation of the model, and hence generates exact draws for any specified exit policy.<sup>1</sup>

## 2 Preliminaries

In this section we briefly review CFTP, and recall the major features of Hopenhayn’s (1992) model.

### 2.1 Coupling From the Past

There are many versions of the CFTP algorithm. We now review a version for finite state Markov chains that provides a useful stepping stone to understanding the algorithm developed below. To begin, consider an irreducible aperiodic Markov chain  $\{X_t\}$  on finite state space  $S$  with unique stationary distribution  $\mu$ . Suppose that the dynamics of the model are given by  $X_{t+1} = h(X_t, W_{t+1})$  where  $\{W_t\}$  is IID. If we start at any  $X_0$  and simulate  $\{X_t\}$  by drawing  $W_1, W_2, \dots$  and successively applying the maps  $h(\cdot, W_1), h(\cdot, W_2), \dots$ , then the distribution of  $X_t$  will converge to (but in general never reach) the stationary distribution  $\mu$ . A second immediate observation is that if  $\{X_t\}$  and  $\{X'_t\}$  are two chains simulated using the *same sequence* of shocks  $\{W_t\}$ , and if  $X_\tau = X'_\tau$  for some  $\tau$ , then  $X_t = X'_t$  for all  $t \geq \tau$ . We say that  $\{X_t\}$  and  $\{X'_t\}$  *couple* at time  $\tau$ .

Now consider an IID sequence of shocks  $\{W_t\}_{t=-\infty}^0$  indexed on  $\{\dots, -2, -1, 0\}$ , and let  $\{X_t\}_{t=-\infty}^0$  be the process generated from these shocks, starting in the infinite past and terminating at  $t = 0$ . Intuitively, since the burn-in is infinite, the distribution of  $X_0$  will be exactly  $\mu$ . On the other hand, because the chain  $\{X_t\}_{t=-\infty}^0$  is infinite and hence cannot be simulated, it appears that  $X_0$  is not observable. In fact this is not the case. To understand how  $X_0$  can be observed, suppose that we fix  $T \in \mathbb{N}$ , take the finite subset of shocks  $W_{-T+1}, W_{-T+2}, \dots, W_0$  and then, for each point  $x$  in the state space  $S$ , construct one version of the process  $\{X_t(x)\}_{t=-T}^0$  that starts at this point  $x$  at time  $-T$  and runs up

---

<sup>1</sup>Exactness is modulo the errors associated with floating point arithmetic, which cannot be avoided.




Figure 1: Coalescence of the tracking processes

until time zero. All of these processes  $\{X_t(x)\}_{t=-T}^0$  are updated using the same shocks  $W_{-T+1}, W_{-T+2}, \dots, W_0$ . We will call them *tracking processes*. Because there are only finitely many tracking processes and each is of finite length, it is possible to simulate them. Two different visualizations of the tracking processes are shown in the top panel of figures 1 and 2 respectively. Here the state space is  $\{1, \dots, 10\}$ , and the two figures show different realizations from the same model. Notice that in the simulations, some of the processes couple and then run together up until time zero. In figure 1, all of the tracking processes have coupled by time zero, and only one final value is observed. If this is the case we will say that the tracking processes *coalesce*. In figure 2 multiple paths are still present at time zero, and the processes fail to coalesce.

Now consider again the process  $\{X_t\}_{t=-\infty}^0$  generated from the entire sequence of shocks  $\{W_t\}_{t=-\infty}^0$ . Since this process must pass through one point  $x$  in the state space at  $-T$ , and since it receives the same shocks  $W_{-T+1}, W_{-T+2}, \dots, W_0$  as the tracking processes from that time forwards, over the period  $-T, -T+1, \dots, 0$  it must follow the same path as the tracking process  $\{X_t(x)\}_{t=-T}^0$  that started at  $x$ . Of course we do not know which of the tracking processes it is following. However, *if* the tracking processes coalesce, then there is only *one* final observation. This observation must be a realization of the time zero value  $X_0$  of the process  $\{X_t\}_{t=-\infty}^0$  that started in the infinite past, and is therefore a draw from the stationary distribution  $\mu$ . Such an outcome is illustrated in the bottom panel of figure 1. For comparison, an unsuccessful outcome is illustrated in the bottom panel of figure 2. Here there are three final values, and we do not know which is the time zero value of the process  $\{X_t\}_{t=-\infty}^0$ . In this case we can take one step further back in time, drawing the additional shock  $W_{-T}$  while preserving the existing shocks  $W_{-T+1}, W_{-T+2}, \dots, W_0$ , recalculate the tracking processes, test for coalescence, and so on. This procedure will




Figure 2: The tracking processes failing to coalesce

eventually terminate with an exact draw from  $\mu$ .<sup>2</sup>

While this technique works when there are a finite number of states, it is clearly not possible to directly implement the same idea when the state space is infinite. However, variations on the idea can potentially be found by exploiting the structure of a given model. In the case of the entry-exit model studied below, we show how the monotonicity of productivity for incumbent firms and the renewal caused by new entrants can be combined to pin down the final value  $X_0$ .

## 2.2 The Model

Hopenhayn (1992) developed what is now a benchmark model for studying the evolution of a competitive industry where entry and exit is endogenously determined. In this section we briefly outline the model and review standard (non-exact) simulation. Most of our notation follows Hopenhayn (1992). If  $X$  is a random variable, then  $\mathcal{D}(X)$  denotes the distribution of  $X$ . The symbol  $\mathcal{B}$  represents the Borel sets on  $[0, 1]$ .

In the model there is a continuum of firms that produce a homogeneous good. The firms face idiosyncratic productivity shocks that follow a Markov process on  $S := [0, 1]$ . The conditional cumulative distribution function for the shock process is denoted by  $F(\phi' | \phi)$ . Following Hopenhayn (1992, assumptions A.3 and A.4), we impose the following restrictions:

---

<sup>2</sup>Using finiteness of the state space, aperiodicity and irreducibility, it can be shown that termination occurs in finite time with probability one.

**Assumption 2.1.**  $F$  is decreasing in its second argument and, for any  $\epsilon > 0$  and any  $\phi \in S$ , there exists an integer  $n$  such that  $F^n(\epsilon | \phi) > 0$ .<sup>3</sup>

We let  $P$  denote the corresponding stochastic kernel. That is,  $P(\phi, A) := \int_A F(d\phi' | \phi)$  for  $\phi \in S$  and  $A \in \mathcal{B}$ . Incumbent firms exit the industry whenever their current productivity falls below a reservation value  $x_t$ . Letting  $M_t$  be the mass of entrants at time  $t$  and  $\nu$  be the Borel probability measure from which the productivity of entrants is drawn, the sequence of firm distributions  $\{\mu_t\}$  on  $S$  satisfies  $\mu_{t+1}(A) = \int P(\phi, A) \mathbb{1}\{\phi \geq x_t\} \mu_t(d\phi) + M_{t+1} \nu(A)$  for all  $A \in \mathcal{B}$ , where  $\mathbb{1}\{\cdot\}$  is an indicator function. At the stationary equilibrium, both  $x$  and  $M$  are constant, and a stationary distribution  $\mu$  is a Borel probability<sup>4</sup> measure  $\mu$  satisfying

$$\mu(A) = \int P(\phi, A) \mathbb{1}\{\phi \geq x\} \mu(d\phi) + M \nu(A) \quad (A \in \mathcal{B}). \quad (1)$$

This is eq. (12) in Hopenhayn (1992). It follows from (1) and  $\mu(S) = P(\phi, S) = \nu(S) = 1$  that  $M = M(x, \mu) := \mu\{\phi \in S : \phi < x\}$ . As a result, we can also write (1) as

$$\mu(A) = \int Q(\phi, A) \mu(d\phi) \quad (2)$$

where

$$Q(\phi, A) := P(\phi, A) \mathbb{1}\{\phi \geq x\} + \nu(A) \mathbb{1}\{\phi < x\}. \quad (3)$$

Equation (2) states that  $\mu$  is a stationary distribution for the stochastic kernel  $Q$  in the usual sense of time invariance. As shown by Hopenhayn (1992, p. 13), the kernel  $Q$  has only one stationary distribution. For the purposes of this paper we will treat  $x$  as given.

## 2.3 Simulation

Our initial task is to produce an ergodic Markov process the stationary distribution of which coincides with the distribution  $\mu$  in (2), taking  $x$  as a given constant in  $(0, 1)$ . To do this, we need a method for sampling from the stochastic kernel  $Q$ . The first step is to simulate from the conditional distribution  $P(\phi, \cdot) = F(\cdot | \phi)$ . In particular, we seek a random variable  $U$  and a function  $g$  such that  $\mathcal{D}(g(\phi, U)) = F(\cdot | \phi)$  for all  $\phi \in S$ . This can be achieved via the inverse transform method, where  $U$  is uniform on  $[0, 1]$  and  $g(\phi, u) = F^{-1}(u | \phi)$ .<sup>5</sup> Now consider the process  $\{\Phi_t\}$  defined by

$$\Phi_{t+1} = g(\Phi_t, U_{t+1}) \mathbb{1}\{\Phi_t \geq x\} + Z_{t+1} \mathbb{1}\{\Phi_t < x\} \quad (4)$$

where  $\{(U_t, Z_t)\}$  is IID with  $\mathcal{D}(Z_t) = \nu$  and  $\mathcal{D}(U_t) = \text{Uniform}[0, 1]$ . Comparing (3) and (4), it can be seen that  $\{\Phi_t\}$  is a Markov process with stochastic kernel  $Q$ .

<sup>3</sup> $F^n(\cdot | \phi)$  is the conditional distribution for productivity after  $n$  periods, given current productivity  $\phi$ .

<sup>4</sup>We focus only on normalized measures, since other cases are just scalar multiples.

<sup>5</sup>Here  $F^{-1}(\cdot | \phi)$  is the generalized inverse of  $F(\cdot | \phi)$ . That is,  $F^{-1}(u | \phi) := \inf\{z : F(z | \phi) \geq u\}$ .

### 3 Exact Sampling

Let  $\{(U_t, Z_t)\}_{t \in \mathbb{Z}}$  be an infinite sequence of IID shocks indexed on  $\mathbb{Z}$  and with each pair  $(U_t, Z_t)$  having the product distribution  $\text{Uniform}[0, 1] \times \nu$ . To simplify notation we will let  $g_t := g(\cdot, U_t)$ , so that, for example,  $g_t \cdots g_1 \phi := g_t \circ g_{t-1} \circ \cdots \circ g_1(\phi)$  is exogenous productivity at  $t$ , given time zero productivity  $\phi \in S$ . To further simplify notation, let

$$h_t(\phi) := g(\phi, U_t) \mathbb{1}\{\phi \geq x\} + Z_t \mathbb{1}\{\phi < x\},$$

so that (4) becomes  $\Phi_{t+1} = h_{t+1} \Phi_t := h_{t+1}(\Phi_t)$ .

In section 2.1 we discussed CFTP for processes with finite state spaces. Since the number of tracking processes is equal to the number of points in the state space, models with finite state space have only finitely many tracking processes. In the current setting, however, the state space  $S$  is a continuum, and hence the number of tracking processes is likewise a continuum. As a result, both occurrence *and* detection of coalescence becomes a considerably more difficult problem.

To describe the tracking processes, fix  $T \geq 1$ . For each  $\phi \in S$ , there is a corresponding tracking process that starts at time  $-T$  with value  $\phi$ , and then updates with maps  $h_{-T+1}, h_{-T+2}, \dots, h_0$ , obtaining the value  $h_0 \cdots h_{-T+1} \phi$  at time zero. We say that the tracking processes *coalesce* if, for some  $T \in \mathbb{N}$ , the set of final values

$$h_0 \cdots h_{-T+1}(S) := \{h_0 \cdots h_{-T+1} \phi : \phi \in S\} \quad (5)$$

is a singleton. What we will now show is that under mild conditions coalescence occurs with probability one, and, moreover, that it is not necessary to keep track of the full continuum of tracking processes in order to find the value of the singleton. In particular, we show that, conditional on a certain event described below, the set of final values  $h_0 \cdots h_{-T+1}(S)$  has only *finitely* many possibilities. Hence coalescence occurs whenever these finite possibilities take the same value. All of these finite possibilities are computable. To begin describing them, let  $T > 1$  be given, and let

$$\Sigma_T := \{k \in \mathbb{N} : 1 \leq k < T \text{ and } g_{-T+k} \cdots g_{-T+2} \cdot g_{-T+1} 1 < x\} \quad \text{and} \quad \sigma_T := \min \Sigma_T.$$

Intuitively,  $\sigma_T$  is the number of periods that an incumbent firm survives, given that it starts at time  $-T$  with maximal productivity 1 and faces the shock sequence  $\{U_t\}_{t \in \mathbb{Z}}$ . Clearly  $\sigma_T$  is only defined when  $\Sigma_T$  is nonempty. However, the probability that  $\Sigma_T$  is nonempty converges to one as  $T \rightarrow \infty$  by assumption 2.1. Moreover, it is remarkable that if  $\Sigma_T$  is nonempty, then the set  $h_0 \cdots h_{-T+1}(S)$ , which contains the final values of the tracking processes started at  $-T$ , can have only finitely many values:

**Lemma 3.1.** *If  $\Sigma_T$  is nonempty, then  $h_0 \cdots h_{-T+1}(S) \subset \Lambda_T$ , where*

$$\Lambda_T := \{h_0 \cdots h_{-T+k+1} Z_{-T+k} : k = 1, \dots, \sigma_T + 1\}. \quad (6)$$




Figure 3: Tracking processes with  $T = 10$  and  $\sigma_T = 8$

The proof of lemma 3.1 is given in section 5. The intuition for the result can be obtained by considering figure 3. In the figure,  $T = 10$ . Tracking processes are plotted for 50 different initial values of  $\phi \in S$ . (Ideally, tracking processes would be plotted from every  $\phi \in S$ , but this is clearly impossible.) For this particular realization of shocks, the set  $\Sigma_T$  is nonempty because the process starting from 1 at time  $-10$  falls below  $x$  at  $t = -2$  (and hence  $\sigma_T = 10 - 2 = 8$ ). As is clear from the figure, the fact that the process starting from 1 at time  $-10$  falls below  $x$  at  $t = -2$  implies that *all* tracking processes fall below  $x$  at least once between  $-10$  and  $-2$  (recall that the productivity of incumbents is monotone). Moreover, if any collection of tracking processes fall below  $x$  at some point in time  $t$ , they subsequently couple, taking the common value  $Z_{t+1}$  at  $t + 1$  and being equal from then on. As a result, by  $t = -1$ , there are at most  $\sigma_T + 1 = 9$  distinct tracking processes. Their time zero values are included in the set  $\Lambda_T$  defined in lemma 3.1. In particular,  $\Lambda_T$  is the time zero values of the processes that start below  $x$  at dates  $-10, -9, \dots, -2$ .

To see the importance of lemma 3.1, let  $\{\Phi_t\}_{t \in \mathbb{Z}}$  be a stationary, doubly-indexed process on the same probability space as  $\{(U_t, Z_t)\}_{t \in \mathbb{Z}}$  that obeys  $\Phi_{t+1} = h_{t+1} \Phi_t$  for all  $t \in \mathbb{Z}$ . The common marginal distribution of  $\Phi_t$  is  $\mu$ . Since  $\Phi_{-T}$  lies somewhere in  $S$ , we know that  $\Phi_0 = h_0 \cdots h_{-T+1} \Phi_{-T} \in h_0 \cdots h_{-T+1}(S)$ . Moreover, if the set  $\Sigma_T$  is nonempty, then lemma 3.1 yields the inclusion  $h_0 \cdots h_{-T+1}(S) \subset \Lambda_T$ , and  $\Phi_0$  lies in the finite observable set  $\Lambda_T$ . In particular, if  $\Lambda_T$  is a singleton, then the value of  $\Phi_0$  is revealed as the value of that singleton.

Figures 4 and 5 show simulations with successful and unsuccessful coalescence respectively. In each figure, the top panel shows only the tracking processes. (As with figure 3, the full continuum of tracking processes cannot be plotted, so we show only 50.) The bottom panel shows the tracking processes and the path of  $\{\Phi_t\}$ . In reality, the path of




Figure 4: Successful coalescence from  $T = 50$




Figure 5: Failure of coalescence

$\{\Phi_t\}$  is not observed. However, in figure 4, there is only one final, coalesced value, and  $\Phi_0$  must take this value. Hence  $\Phi_0$  is observed. On the other hand, in figure 5,  $\Phi_0$  is equal to one of two final values, and we have no way of identifying which one it is.

Now let us consider how to use our results to sample from  $\mu$  by generating observations of  $\Phi_0$ . In order to avoid conditioning on coalescence by a certain point in time, we wish to reveal the value of  $\Phi_0$  for every random seed. This can be done by fixing the seed, which determines the values of the shock processes, and then taking  $T$  larger and larger until coalescence occurs. Algorithm 1 gives details. The algorithm terminates with an exact draw from  $\mu$ . Replication with independent shocks will generate independent draws.

---

**Algorithm 1:** Generates an exact draw from  $\mu$

---

```

fix  $T$  to be an integer greater than 1;
draw  $(U_0, Z_0), \dots, (U_{-T+1}, Z_{-T+1})$  independently from their correct distributions;
repeat
  compute the set  $\Sigma_T$  ;
  if  $\Sigma_T$  is nonempty then
    compute the set  $\Lambda_T$  ;
    if  $\Lambda_T$  is a singleton then
      set  $\Phi_0$  to be the value of that singleton ;
      break ;
    end
  end
  draw  $(U_{-T}, Z_{-T})$  and append to the end of list  $(U_0, Z_0), \dots, (U_{-T+1}, Z_{-T+1})$  ;
  set  $T = T + 1$  ;
end
return  $\Phi_0$  ;

```

---

At this stage we do not know that the algorithm will terminate with probability one. This issue is central to the correctness of the algorithm because, as discussed above, the way we avoid conditioning is by revealing the value of  $\Phi_0$  for every random seed. We now show that probability one termination in finite time holds under the following condition, which is satisfied by many standard distributions.

**Assumption 3.1.** If  $G \subset S$  is a nonempty open set, then  $\nu(G) > 0$ .

**Proposition 3.1.** Let  $T^* := \min\{T \in \mathbb{N} : \Sigma_T \text{ is nonempty and } \Lambda_T \text{ is a singleton}\}$ . If assumption 3.1 holds, then  $\mathbb{P}\{T^* < \infty\} = 1$ .

The proof of proposition 3.1 is given in section 5. A simple implementation of algorithm 1 can be obtained from [http://johnstachurski.net/papers/hh\\_sampling.html](http://johnstachurski.net/papers/hh_sampling.html). We tested the code with two different parameterizations. First, we followed Hopenhayn




Figure 6: 95% confidence set for the stationary distribution




Figure 7: Stationary density, first parameterization

and Rogerson (1993) in taking the distribution  $\nu$  for new entrants to be uniform, and the process for incumbents to be  $\Phi_{t+1} = g_{t+1}\Phi_t = a + \rho\Phi_t + \epsilon_{t+1}$  where  $\{\epsilon_t\}$  is IID with distribution  $N(0, \sigma^2)$ .<sup>6</sup> The parameters were set to  $a = 0.36$ ,  $\rho = 0.4$  and  $\sigma = 0.1$ , while  $x$  was set to 0.49, so that approximately 40% of incumbents exit within 5 years (see Hopenhayn, 1992, p. 1127). For these parameters, running the program on our computer produces about 36,000 independent draws from  $\mu$  per second.<sup>7</sup>

Figure 6 shows a 95% confidence set for the cumulative distribution function corresponding to  $\mu$ , based on 36,000 observations and calculated using the Kolmogorov distribution of the sup norm deviation between true and empirical cdfs. The Kolmogorov result is applicable here because the draws are exact and IID. The true distribution function lies entirely between the two bands with 95% probability. Figure 7 shows the corresponding

<sup>6</sup>To bound the process we added reflecting barriers at 0 and 1.

<sup>7</sup>The computer is a standard workstation with a 2.67GHz Intel CPU and 4 gigabytes of RAM.




Figure 8: Stationary density, second parameterization

density, computed from the same observations combined with a standard nonparametric kernel density estimator (using a Gaussian kernel).

For the second parameterization we took the process for incumbents to be  $\Phi_{t+1} = \Phi_t B_{t+1}$ , where  $B_t \sim \text{Beta}(5, 1)$ , and set  $\nu = \text{Beta}(5, 1)$ . We set  $x$  to 0.35, so that approximately 40% of firms exit within 5 years. For these parameters, the program produces about 30,000 independent draws from  $\mu$  per second. Figure 8 shows the stationary density, computed from 30,000 observations and a standard nonparametric kernel density estimator.

As a final illustration of the algorithm, suppose now that we wish to calculate aggregate output  $\int f(\phi, \ell) \mu(d\phi)$ , where  $f$  is a production function and  $\ell$  is given. Letting  $f(\phi, \ell) = \phi \ell^\theta$  with  $\ell = 0.5$  and  $\theta = 0.64$ , and adopting the first set of parameters, we computed the empirical estimate  $Q_n := \frac{1}{n} \sum_{i=1}^n f(\Phi_i, \ell)$  using 36,000 observations (generated in one second). The mean  $Q_n$  was 0.3848, while the standard error computed from the CLT was 0.00085, giving a 95% confidence interval for  $\int f(\phi, \ell) \mu(d\phi)$  of  $[0.3840, 0.3857]$ . With more draws the standard errors fall at rate  $O(n^{-1/2})$ .

## 4 Conclusion

In this paper we developed an algorithm that generates exact draws from the stationary distribution of Hopenhayn's (1992) entry-exit model for a given exit policy. In the applications we considered, the algorithm produced in the order of 30,000 to 36,000 independent exact draws from the stationary distribution per second. The basic idea should extend to many economic applications involving entry and exit with monotone exogenous shocks.

## 5 Proofs

In the following proofs we write  $\sigma$  for  $\sigma_T$  to simplify notation.

*Proof of lemma 3.1.* Let  $\Sigma_T$  be nonempty. As a first step, we show that if  $\phi \geq x$ , then there exists a  $j \in \{1, \dots, \sigma\}$  such that  $h_{-T+j} \cdots h_{-T+1} \phi < x$ . To see that this is so, fix  $\phi \geq x$  and suppose that the statement fails. In other words,  $h_{-T+j} \cdots h_{-T+1} \phi \geq x$  for  $j \in \{1, \dots, \sigma\}$ . We know that if  $y \geq x$ , then  $h_i y = g_i y$ . It follows that  $h_{-T+\sigma} \cdots h_{-T+1} \phi = g_{-T+\sigma} \cdots g_{-T+1} \phi$ . But then

$$x \leq h_{-T+\sigma} \cdots h_{-T+1} \phi = g_{-T+\sigma} \cdots g_{-T+1} \phi \leq g_{-T+\sigma} \cdots g_{-T+1} 1 < x,$$

where the second inequality is due to monotonicity of  $g_i$ , and then third is by the definition of  $\sigma$ . Contradiction.

To complete the proof, pick any  $\phi \in S$ . Our claim is that  $h_0 \cdots h_{-T+1} \phi \in \Lambda_T$ . Suppose first that  $\phi < x$ . In this case we have  $h_0 \cdots h_{-T+1} \phi = h_0 \cdots h_{-T+2} Z_{-T+1}$ , which is an element of  $\Lambda_T$ . Next, suppose that  $\phi \geq x$ . In light of the preceding argument, there exists a  $j \in \{0, \dots, \sigma\}$  with  $h_{-T+j} \cdots h_{-T+1} \phi < x$ , and hence

$$h_{-T+j+1} \cdot h_{-T+j} \cdots h_{-T+1} \phi = Z_{-T+j+1},$$

from which we obtain

$$\begin{aligned} h_0 \cdots h_{-T+1} \phi &= h_0 \cdots h_{-T+j+2} \cdot h_{-T+j+1} \cdot h_{-T+j} \cdots h_{-T+1} \phi \\ &= h_0 \cdots h_{-T+j+2} Z_{-T+j+1}. \end{aligned}$$

Since  $j \in \{0, \dots, \sigma\}$ , the right-hand side is an element of  $\Lambda_T$ . This completes the proof.  $\square$

*Proof of proposition 3.1.* Let  $n$  be an integer such that  $F^n(x | 1) > 0$ , existence of which is due to assumption 2.1. Fixing  $j \in \mathbb{N}$ , let

$$E_j := \{g_{-(j-1)n-1} \cdots g_{-jn} 1 < x\} \cap \{Z_{-(j-1)n-1} < x, \dots, Z_{-jn} < x\}.$$

The events  $\{g_{-(j-1)n-1} \cdots g_{-jn} 1 < x\}$  and  $\{Z_{-(j-1)n-1} < x, \dots, Z_{-jn} < x\}$  are independent because the first event depends only on  $U_{-(j-1)n-1}, \dots, U_{-jn}$  and the second depends only on  $Z_{-(j-1)n-1}, \dots, Z_{-jn}$ . As a result,

$$\delta := \mathbb{P}(E_j) = F^n(x | 1) \nu([0, x])^n.$$

The constant  $\delta$  is strictly positive as a result of assumption 3.1. We claim that if the event  $E_j$  occurs, then  $\Sigma_{jn+1}$  is nonempty and  $\Lambda_{jn+1}$  is a singleton. To simplify notation, we treat only the case of  $j = 1$ .

So suppose that  $E_1$  occurs. Clearly  $\Sigma_{n+1}$  contains  $n$ , and hence is nonempty. To see that  $\Lambda_{n+1}$  is a singleton, observe that since  $\sigma = \sigma_{n+1}$  is the smallest element of  $\Sigma_{n+1}$ , we must have  $\sigma \leq n$ . As a consequence,

$$\begin{aligned}\Lambda_{n+1} &= \{h_0 \cdots h_{-n+k} Z_{-n-1+k} : k = 1, \dots, \sigma + 1\} \\ &\subset \{h_0 \cdots h_{-n+k} Z_{-n-1+k} : k = 1, \dots, n + 1\}.\end{aligned}$$

We claim that on the set  $E_1$  we have

$$h_0 \cdots h_{-n+k} Z_{-n+k-1} = Z_0 \quad \text{for any } k \in \{1, \dots, n + 1\}. \quad (7)$$

To prove that (7) holds, observe that on  $E_1$  the values  $Z_{-1}, \dots, Z_{-n}$  are all less than  $x$ . As a result, we have

$$\begin{aligned}h_0 Z_{-1} &= Z_0 \\ h_0 h_{-1} Z_{-2} &= h_0 Z_{-1} = Z_0 \\ h_0 h_{-1} h_{-2} Z_{-3} &= h_0 h_{-1} Z_{-2} = h_0 Z_{-1} = Z_0\end{aligned}$$

and so on. Together, these equalities give (7). As a consequence, we conclude that  $\Sigma_{n+1}$  is nonempty and  $\Lambda_{n+1}$  is a singleton whenever  $E_1$  occurs, and, more generally,  $\Sigma_{nj+1}$  is nonempty and  $\Lambda_{nj+1}$  is a singleton whenever  $E_j$  occurs. The events  $E_1, E_2, \dots$  are independent and have positive probability  $\delta$ . As a consequence, the set  $\cup_j E_j$  has probability one. In other words, with probability one, there exists a finite  $T \in \mathbb{N}$  such that  $\Sigma_T$  is nonempty and  $\Lambda_T$  is a singleton.  $\square$

## References

- [1] Athreya, K. B., Stenflo, O., 2003. Perfect sampling for Doeblin chains. *Sankhya* 65, 763–777.
- [2] Clementi, G. L. and B. Palazzo, 2010. Entry, exit, firm dynamics, and aggregate fluctuations. Mimeo.
- [3] Cooley, T.F. and Quadrini, V., 2001. Financial markets and firm dynamics. *American Economic Review*, 91 (5), 1286–1310.
- [4] Corcoran, J. N., Tweedie, R. L., 2001. Perfect sampling of ergodic Harris chains. *Annals of Applied Probability* 11, 438–451.
- [5] Foss, S., Tweedie, R. L., 1998. Perfect simulation and backward coupling. *Stochastic Models* 14, 187–204.
- [6] Hopenhayn, H. A., 1992. Entry, exit and firm dynamics in long run equilibrium. *Econometrica*, 60 (5), 1127–1150.

- [7] Hopenhayn, H. and Rogerson, R., 1993. Job turnover and policy evaluation: A general equilibrium analysis. *Journal of Political Economy*, 101 (5), 915–938.
- [8] Levin, D. A., Y. Peres, E. L. Wilmer, 2008. *Markov Chains and Mixing Times*. American Mathematical Society.
- [9] Melitz, M.J., 2003. The impact of trade on intra-industry reallocations and aggregate industry productivity. *Econometrica*, 71 (6), 1695–1725.
- [10] Meyn, S., Tweedie, R. L., 1993. *Markov Chains and Stochastic Stability*. Springer-Verlag, New York.
- [11] Murdoch, D. J., Green, P. J., 1998. Exact sampling from a continuous state space. *Scandinavian Journal of Statistics* 25, 483–502.
- [12] Nishimura, K. and J. Stachurski, 2010. Perfect simulation of stationary equilibria. *Journal of Economic Dynamics and Control*, 34, 577–584.
- [13] Propp, J. G., Wilson, D. B., 1996. Exact sampling with coupled Markov chains and applications to statistical mechanics. *Random Structures and Algorithms* 9, 223–252.